PROSPECTS
IN PHARMACEUTICAL SCIENCES

Prospects in Pharmaceutical Sciences, 23(2), 67-73
https://prospects.wum.edu.pl/

Original Article

HERBAL POWER: ANTIMICROBIAL, ANTIOXIDANT AND ANTICHOLINESTERASE PROPERTIES OF *CITRULLUS COLOCYNTHIS*

Emre Sevindik[1], Orhan Unal[2], Imran Uysal[3], Mansur Seymen Seğmenoğlu[4], Mustafa Sevindik*[5]

[1] *Department of Agricultural Biotechnology, Faculty of Agriculture, Aydın Adnan Menderes University, 09-100 Aydın, Türkiye*

[2] *Department of Biology, Faculty of Science, Akdeniz University, 07-070 Antalya, Türkiye.*

[3] *Department of Food Processing, Bahçe Vocational School, Osmaniye Korkut Ata University, 80-500 Osmaniye, Türkiye.*

[4] *Department of Nursing, Faculty of Health Sciences, Osmaniye Korkut Ata University, 80-000 Osmaniye, Turkey*

[5] *Department of Biology, Faculty of Engineering and Natural Sciences, Osmaniye Korkut Ata University, 80-000 Osmaniye, Türkiye.*

* Correspondence, e-mail: sevindik27@gmail.com

Received: 6.02.2025 / Accepted: 20.02.2025 / Published: xx.05.2025

ABSTRACT

Citrullus colocynthis (L.) Schrad. is a biologically rich plant widely used in traditional medicine. This study aimed to investigate the antimicrobial, antioxidant and anticholinesterase potential and phenolic/flavonoid contents of *C. colocynthis*. Antimicrobial tests using ethanol and methanol extracts revealed that it exhibited strong antimicrobial effects especially against pathogens such as *Staphylococcus aureus*, *Candida albicans*, *C. krusei* and *Acinetobacter baumannii*. In addition, antioxidant activity tests showed that ethanol extract had high antioxidant capacity, while methanol extract was moderately effective. Phenolic and flavonoid content analyses showed that ethanol extract contained higher phenolic and flavonoid components. In addition, anticholinesterase activity tests showed that *C. colocynthis* extract exhibited a significant anticholinesterase effect, which could potentially be useful in the treatment of neurological diseases. These findings indicate that *C. colocynthis* is a potent natural resource with potential therapeutic benefits and its antimicrobial, antioxidant and anticholinesterase properties are of considerable importance in biomedical applications.

KEYWORDS: Antialzeihmer, Antimicrobial, Antioxidant, *Citrullus colocynthis*, Medicinal plants.

Article is published under the CC BY license.

1. Introduction

Since ancient times, people have relied on natural products obtained from plants against various diseases [1]. These medicinal plants, one of the cornerstones of traditional medicine, are rich in bioactive substances and attract attention with their therapeutic effects [2]. At the same time, they are an important natural source for the discovery of new drugs used in the treatment of diseases in modern medicine [3,4]. Scientific studies have revealed that different plant species have many biological properties such as antimicrobial, antioxidant, anticancer, antiproliferative, hepatoprotective, antiaging, antiallergic and DNA protective [5-9]. These properties support the versatile use potential of herbal ingredients in the field of health.

Citrullus colocynthis (L.) Schrad. is a medicinally important plant belonging to the Cucurbitaceae family [10]. This plant, which grows widely in desert regions, has a high nutraceutical potential and is widely used in medical and pharmaceutical fields [11]. Known as "bitter apple" among the public, this plant is also known by different names such as colocynth, bitter cucumber, bitter melon, egusi melon and bitter vine of Sodom [12,13]. *C. colocynthis*, which has been used in traditional medicine for many years, has been reported to be effective in the treatment of various diseases such as cold, cough, bronchitis, asthma, diabetes, jaundice, dysentery and cancer [13]. This plant, which has been evaluated for medicinal purposes in different cultures

throughout history, has been reported to have significant potential in the field of health thanks to its bioactive components [13,14].

In this study, antioxidant, antimicrobial, antiallergic, total phenolic and total flavonoid contents of *C. colocynthis* were determined.

2. Materials and Methods

C. colocynthis plant samples were collected from Duhok region of Iraq. Collected plant samples are kept in the herbarium of Biology Department of Faculty of Science, Zakho University. For experimental studies, plant samples were first dried and powdered. Then, 30 grams of prepared powder samples were weighed and extracted with 250 mL of ethanol in Soxhlet apparatus at 50 °C for approximately 6 hours. After extraction, solvents were removed using rotary evaporator and the obtained extract was stored at +4 °C until experiments were performed.

2.1. Antimicrobial activity tests

In this study, antibacterial and antifungal activities of *C. colocynthis* plant extract were determined using agar dilution method. Bacterial strains used in the study were *Staphylococcus aureus* ATCC 29213, *S. aureus* MRSA ATCC 43300, *Enterococcus faecalis* ATCC 29212, *Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853 and *Acinetobacter baumannii* ATCC 19606. Fungal strains used were *Candida albicans* ATCC 10231, *C. krusei* ATCC 34135 and *C. glabrata* ATCC 90030. Bacterial strains were precultured in Müller Hinton Broth and fungal strains were precultured in RPMI 1640 Broth before the experiment. The plant extract was tested by preparing stock solutions in the concentration range of 12.5-800 µg/mL. The lowest concentrations that inhibited the growth of microorganisms were determined and the results obtained were reported in µg/mL [15-18].

2.2. Antioxidant tests

Total antioxidant (TAS) and total oxidant (TOS) levels of *C. colocynthis* plant extract were determined using Rel Assay kits. The protocol provided by the manufacturer of the kits was followed in the analyses. TAS values were expressed as mmol Trolox equivalent/L, while TOS values were expressed as µmol hydrogen peroxide equivalent/L [19,20]. Oxidative stress index (OSI) is a parameter that shows the extent to which oxidant compounds are suppressed by antioxidant compounds. In the OSI calculation, firstly the units of TOS and TAS values were equalized, and then the ratio of the TOS value to the TAS value was taken and expressed as a percentage [21].

2.3. Anticholinesterase tests

Anticholinesterase activity of ethanol extract of *C. colocynthis* plant was determined using Ellman method [22]. In the study, plant extract was prepared at different concentrations in the range of 200-3.125 µg/mL using distilled water. For analysis, 130 µL of 0.1 M (pH=8) phosphate buffer, 10 µL of stock solution and 20 µL of enzyme solution (acetylcholinesteras (AchE) or butyrylcholinesterase (BChE)) were added to the microplate wells, respectively. The mixture was incubated at 25 °C in the dark for 10 min. After the incubation period, 20 µL of DTNB (5,5'-dithiobis-(2-nitrobenzoic acid)) solution and 20 µL of substrate (acetylcholine iodide or butyrylcholine iodide) were added to each well and the absorbance of the reaction was measured at 412 nm wavelength. Experiments were performed in three parallel repetitions and the results obtained were expressed in µg/mL by calculating IC50 values.

2.4. Total Phenolic and Total Flavonoid Analyses

A 1 mL stock solution was prepared from the ethanol extract of *C. colocynthis* plant. In order to determine the amount of phenolic compounds, 1 mL of Folin-Ciocalteu reagent (1:9, v/v) was added to the prepared solution and mixed. Then, 0.75 mL of 1% Na_2CO_3 was added to the mixture and incubated for 2 hours at room temperature. At the end of the incubation period, the absorbance value was measured at 760 nm and the total phenolic content (TPC) was calculated in mg/g using the calibration curve of the gallic acid standard solution [23].

The total flavonoid content (TFC) of the ethanol extract of the plant was determined using the aluminum chloride test. For the analysis, 0.1 mL of 10% $Al(NO_3)_3$, 0.1 mL of 1 M NH_4CH_3COO, 4.3 mL of methanol, 0.5 mL of quercetin and 0.5 mL of plant extract were mixed. After the obtained solution was incubated for approximately 40 minutes, the absorbance value was measured at 415 nm. The total flavonoid content was calculated and expressed in mg/g [23].

3. Results and Discussion

3.1. Antimicrobial activity

Today, the origin of many diseases is microorganisms. The effects of drugs used to combat microorganisms are insufficient due to resistant microorganisms. In this context, the discovery of new antimicrobial drugs has become inevitable [24]. Many medicinal plants show strong antimicrobial activity against various bacterial and fungal species thanks to the bioactive components they contain, and these properties make them an important source for natural treatment alternatives and new drug discoveries [25,26]. In our study, the lowest extract concentration of *C. colocynthis* that prevented the growth of test microorganisms was determined. The findings are shown in Table 1.

Table 1. Antimicrobial potential of *Citrullus colocynthis*

Microorganisms	Solvent	
	Ethanol	Methanol
S. aureus	50	100
S. aureus MRSA	50	100
E. faecalis	100	200
E. coli	100	100
P. aeruginosa	200	400
A. baumannii	50	50
C. glabrata	100	100
C. albicans	50	100
C. krusei	50	100

*50, 100, 200 and 400 µg/mL are the lowest extract concentrations that inhibit the growth of microorganisms.

In the literature, *C. colocynthis* is reported to be infected with *Enterococcus faecalis, Staphylococcus*

aureus, S. epidermidis, Listeria monocytogenes, Micrococcus luteus, Escherichia coli, Pseudomonas aeruginosa, Salmonella typhimurium, Vibrio parahaemolyticus, Vibrio alginolyticus, Candida albicans, Candida glabrata, Candida kreusei and *Candida parapsilosis* [27]. In a different study, it was reported that aqueous and methanol extracts of *C. colocynthis* were effective against *Bacillus subtilis, Staphylococcus aureus, Streptococcus faecalis, Streptococcus pyogenes, Escherichia coli, Klebseilla pneumoniae, Proteus mirabilis, Salmonella typhi, Vibrio cholerae, Aspergillus fumigatus, Mucor* sp. and *Penicillium* sp. [28]. In our study, ethanol and methanol extracts of *C. colocynthis* were used. As a result of the study, it was determined that ethanol extracts generally exhibited stronger antimicrobial activity compared to methanol extracts. Ethanol extract at 50 µg/mL and methanol extract at 100 µg/mL were effective against *S. aureus, S. aureus* MRSA, *C. albicans* and *C. krusei*. Both ethanol and methanol extracts at 100 µg/mL were effective against *E. coli* and *C. glabrata*. Ethanol extract at 100 µg/mL and methanol extract at 200 µg/mL were effective against *E. faecalis*. Ethanol extract at 200 µg/mL and methanol extract at 400 µg/mL were effective against *P. aeruginosa*. Both ethanol and methanol extract at 50 µg/mL were effective against *A. baumannii*. In this context, it was determined in our study that *C. colocynthis* was effective at concentrations of 50-400 µg/mL. As a result, it was determined that *C. colocynthis* has antimicrobial potential. This antimicrobial effect is due to the bioactive components of the plant, and it is known that compounds such as flavonoids, alkaloids, terpenes and phenolic acids damage the cell walls of pathogenic microorganisms and inhibit their biosynthetic pathways [29,30]. In addition, some studies have reported that the antibacterial and antifungal effects of *C. colocynthis* are also effective on microorganisms resistant to traditional treatment methods, which increases the importance of the plant as a potential source in the pharmaceutical field. Similarly, in our study, it was determined that ethanol and methanol extracts were effective against important pathogens, especially *S. aureus, Candida albicans* and *Pseudomonas aeruginosa*. These findings reveal the potential of *C. colocynthis* as a natural treatment alternative that can be used in the treatment of resistant infections in modern medicine.

3.2. Antioxidant activity

Oxidative stress is a condition that causes cellular damage as a result of the accumulation of free radicals and reactive oxygen species (ROS) in the body. Normally, the body has antioxidant systems to neutralize these harmful components [31]. However, excessive free radical production can have negative effects on cellular structures such as lipid peroxidation, protein denaturation, and DNA damage, which can lead to the development of many diseases such as aging, cancer, cardiovascular diseases, and neurological disorders [32]. Plants exhibit strong antioxidant properties with bioactive components such as phenolic compounds, flavonoids, alkaloids, and terpenoids [33]. These compounds can help protect cells from oxidative damage by neutralizing free radicals [34]. Plant antioxidants, in addition to their direct free radical scavenging ability, also provide more comprehensive protection by increasing the activity of the body's own antioxidant enzymes [35]. Therefore, the antioxidant potential of plants has become an important area of research in modern medicine in the treatment of diseases caused by oxidative stress [36]. In our study, the antioxidant potential of ethanol and methanol extracts of *C. colocynthis* was determined. The obtained findings are shown in Table 2.

Table 2. TAS, TOS, OSI, TPC and TFC values of *Citrullus colocynthis*

	Ethanol	Methanol
TAS (mmol/L)	8.750±0.147	7.453±0.153
TOS (µmol /L)	10.678±0.083	10.951±0.078
OSI (TOS/(TAS*10))	0.122±0.004	0.147±0.003
TPC (mg/g)	92.74±1.47	86.41±1.38
TFC (mg/g)	149.63±2.37	112.43±2.18

In the literature, the antioxidant potential of *C. colocynthis* has been reported using different methods [37-39]. In our study, TAS, TOS and OSI values were determined using Rel Assay kits. The antioxidant potential of *C. colocynthis* has not been reported with Rel Assay kits in the literature. It was detected for the first time in our study. TAS, TOS and OSI values of different plant species were reported using Rel Assay kits. In this context, TAS values of *Mentha longifolia, Anthemis cotula, Hypericum spectabile, Arum dioscoridis, Glaucium alakirensis* and *Alcea kurdica* were reported as 6.094, 7.625, 9.306, 6.486, 3.496 and 3.298 mmol/L. TOS values were reported as 14.050, 11.247, 13.065, 13.578, 2.204 and 8.312 µmol/L. OSI values were reported as 0.231, 0.148, 0.140, 0.209, 0.063 and 0.252 [40-45].

Compared to these studies, the TAS value of the ethanol extract of *C. colocynthis* was determined to be higher than *M. longifolia, A. cotula, A. dioscoridis, G. alakirensis* and *A. kurdica*, and lower than *H. spectabile*. The TAS value of methanol extract of *C. colocynthis* was determined to be higher than *M. longifolia, A. dioscoridis, G. alakirensis* and *A. kurdica*, and lower than *A. cotula* and *H. spectabile*. The TAS value is an indicator of the totality of antioxidant compounds produced in natural products [46]. The high TAS values of *C. colocynthis* show that this plant is an important antioxidant source and its capacity to neutralize free radicals is significant compared to some other plants. However, when the TAS values of ethanol and methanol extracts were compared with different plants, it was observed that some plants were higher and some plants were lower. This result suggests that the content of antioxidant compounds of *C. colocynthis* may vary from plant to plant and these differences may depend on the extraction method and the properties of the solvent.

The TOS value is an indicator of the totality of oxidant compounds produced in natural products [46]. In our study, it was observed that the TOS values of both ethanol and methanol extracts of *C. colocynthis* were lower than *M. longifolia, A. cotula, H. spectabile* and *A. dioscoridis*, and higher than *G. alakirensis* and *A. kurdica*. Since the TOS value indicates the level of oxidant compounds in plants, a low value suggests that the plant has a balanced structure in terms of oxidant components. In our study, it was determined that the TOS values of ethanol and methanol extracts of *C. colocynthis* were lower than some plants and higher than some plants. This situation reveals that the potential of the plant to produce different oxidant compounds is shaped by

environmental factors and the genetic characteristics of the plant. Regulation of oxidant compounds can strengthen the plant's defense mechanisms, which can contribute to being more resistant to stress factors in the natural environment.

The OSI value shows how much oxidant compounds are suppressed by antioxidant compounds [46]. The OSI value of the ethanol extract of *C. colocynthis* used in our study was determined to be lower than *M. longifolia*, *A. cotula*, *H. spectabile*, *A. dioscoridis* and *A. kurdica*, and higher than *G. alakirensis*. The OSI value of the methanol extract of *C. colocynthis* was determined to be lower than *M. longifolia*, *A. cotula*, *A. dioscoridis*, *A. kurdica*, and higher than *H. spectabile* and *G. alakirensis*. In this context, it was determined that *C. colocynthis* has a significant potential in suppressing oxidant compounds. The OSI value is an important parameter because it shows the interaction of antioxidant and oxidant compounds and the suppression rate of this interaction on the body. In our study, the OSI values of ethanol and methanol extracts of *C. colocynthis* revealed that this plant has an important potential in suppressing oxidant compounds. While the OSI value of the ethanol extract was lower than many plants, it was observed that the methanol extract was lower than certain plants. This suggests that methanol is a more effective solvent and that the methanol extract of *C. colocynthis* may have a stronger oxidative balancing capacity. As a result, it was determined that *C. colocynthis* showed the potential to balance antioxidant and oxidant components with high TAS and appropriate TOS values. These properties suggest that the plant may be an important source that can be used both in preventing oxidative stress and strengthening natural defense mechanisms. However, further research is needed to support the validity of these findings in larger studies and to better understand the clinical and pharmaceutical potential of *C. colocynthis*.

3.3. Total phenolic and total flavonoid contents

Plants are rich in phenolic and flavonoid components, and these components play a role in the defense mechanisms of the plant and may have beneficial effects on health by exhibiting various biological activities [47]. In our study, total phenolic and total flavonoid contents of ethanol and methanol extracts of *C. colocynthis* were determined. The findings are shown in Table 2. As a result of the analyzes, the TPC values of ethanol and methanol extracts of *C. colocynthis* were determined as 92.74±1.47 and 86.41±1.38 mg/g, respectively. In addition, TFC values were determined as 149.63±2.37 and 112.43±2.18 mg/g, respectively. In the literature, the total phenolic content of *C. colocynthis* was reported as 5.39 mg/100 g and the total flavonoid content as 938.0 mg/100 g [48]. Compared to this study, the total phenolic content (TPC) value determined in our study was significantly higher, while the total flavonoid content (TFC) was determined at lower levels. This difference may be due to the differences in the extraction methods used, solvent type, plant growth conditions, developmental stage and analysis techniques. The amount of phenolic and flavonoid components may vary depending on the ecological factors (such as soil structure, water stress, duration of exposure to sunlight) where the plant grows [49]. In addition, the chemical polarity of the solvent used in the extraction method may affect the extraction efficiency of different phenolic compounds [50]. The higher TPC and TFC values of the ethanol extract compared to the methanol extract in our study may be attributed to the fact that ethanol can dissolve flavonoids and phenolic compounds more effectively.

In addition, the fact that the total flavonoid content reported in the literature is quite high, but was measured at lower levels in our study may also be due to differences in the analysis methods used. The aluminum chloride spectrophotometric method used to determine flavonoids may be more sensitive to some flavonoid compounds, and this may lead to variable results among studies [51].

In conclusion, these differences in phenolic and flavonoid content are important in understanding the biological activity and pharmacological potential of the plant. Future studies may provide more comprehensive information by optimizing the extraction parameters, detailed characterization of phenolic and flavonoid components, and investigating their effects on biological activities.

3.4. Anticholinesterase activity

The anticholinesterase activity of plants is important for the discovery of natural compounds that offer a potential treatment options, especially in the treatment of neurological diseases. These activities aim to improve neurological functions by inhibiting the acetylcholinesterase enzyme by the active compounds in herbal extracts and increasing acetylcholine levels [52,53]. In our study, acetylcholinesterase and butyrylcholinesterase activities of *C. colocynthis* were determined. In this context, the potential of the plant against neurodegenerative diseases such as Alzheimer's was determined. IC50 values of the obtained findings are shown in Table 3.

Table 3. Anti-AChE and anti-BChE values of *Citrullus colocynthis*

	AChE (µg/mL)	BChE (µg/mL)
Ethanol	14.96±1.16	31.00±0.96
Methanol	22.62±1.06	41.60±1.09
Galantamine	7.38±0.27	14.33±0.42

It has been reported in the literature that the acetylcholinesterase activity of the methanol extract of *C. colocynthis* is 131.01 µg/mL [54]. Compared to this study, the methanol extract of *C. colocynthis* used in our study exhibited higher acetylcholinesterase activity. There are several possible reasons for these differences. First of all, the biological activity of herbal extracts may vary depending on various such as growing conditions (soil structure, climate, geographical region), harvest time and extraction methods [50]. In addition, different solvents such as ethanol and methanol can dissolve and extract different components in herbal extracts. Ethanol is a solvent generally used to obtain extracts rich in polyphenols and flavonoids, and it is known that these compounds play a role in acetylcholinesterase inhibition [55]. In our study, the higher activity of the ethanol extract compared to the methanol extract suggests that ethanol may have extracted more effective compounds that may act as acetylcholinesterase inhibitors. However, when compared to the standard galantamine, both

extracts were found to have lower activity. This may be due to impurities, low specific activities or insufficiently high inhibitory concentrations of active compounds found in herbal extracts [56]. Therefore, fractionation and isolation studies are needed to make the extracts more effective. Future studies should include detailed analysis of the chemical profiles of the extracts, identification of the main active compounds and elucidation of the mechanisms of their inhibitory effects on acetylcholinesterase. In particular, identification of the phytochemical components underlying the higher activity of the ethanol extract compared to the methanol extract may help us better understand the therapeutic potential of the plant.

4. Conclusion

The findings of this study highlight the significant biological potential of *C. colocynthis* extracts, specifically those obtained through ethanol and methanol. Both extracts exhibited notable antimicrobial and antioxidant activities, with the ethanol extract showing a particularly strong antimicrobial effect. Additionally, both extracts demonstrated substantial acetylcholinesterase inhibitory activity, suggesting a potential role for *C. colocynthis* in the management of neurodegenerative diseases, such as Alzheimer's disease. The observed biological activities correlate with the high concentrations of phenolic compounds and flavonoids present in the extracts, which are likely responsible for their bioactivity. This study provides scientific support for the traditional use of *C. colocynth*is in herbal medicine and underscores its potential as a promising candidate for further pharmaceutical development. However, additional studies are required to evaluate the toxicity profile, pharmacokinetics, and clinical efficacy of *C. colocynthis* extracts before considering their therapeutic application.

Author Contributions: Conceptualization, E.S., O.U., I.U., M.S.S. and M.S.; methodology, E.S., O.U., M.S.S. and M.S.; validation, E.S., I.U., M.S.S. and M.S.; investigation, O.U., I.U., M.S.S. and M.S.; resources, E.S., O.U., I.U. and M.S.; data curation, E.S., O.U., I.U. and M.S.S.; writing—original draft preparation, E.S., O.U., I.U., M.S.S. and M.S.; writing—review and editing, E.S., O.U., I.U., M.S.S. and M.S.; visualization, E.S., O.U., I.U., M.S.S. and M.S.; supervision, E.S., O.U., I.U., M.S.S. and M.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgements

We would like to thank Dr. Falah Saleh Mohammed for his support in procuring the plant.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Karaman, S.; Ilcim, A.; Comlekcioglu, N. Composition of the essential oils of *Salvia aramiensis* Rech. fil. and *Salvia cyanescens* Boiss. & Bal. *Pak. J. Bot.* **2007**, *39(1),* Art. No: 169.
2. Vaou, N.; Stavropoulou, E.; Voidarou, C.; Tsigalou, C.; Bezirtzoglou, E. Towards advances in medicinal plant antimicrobial activity: A review study on challenges and future perspectives. *Microorganisms* **2021**, *9(10),* Art. No: 2041. DOI: 10.3390/microorganisms9102041
3. Kalkan, M.; Aygan, A.; Çömlekçioglu, N.; Çömlekçioğlu, U. *Olea europaea* Yapraklarının Bazı Biyoaktif Özelliklerinin Araştırılması, Antimikrobiyal ve Enzim İnhibisyon Etkinliğinin İncelenmesi. *Turkish Journal of Agriculture-Food Science and Technology* **2023**, *11(3),* 496-504. DOI: 10.24925/turjaf.v11i3.496-504.5828
4. Nwozo, O.S.; Effiong, E.M.; Aja, P.M.; Awuchi, C.G. Antioxidant, phytochemical, and therapeutic properties of medicinal plants: A review. *Int. J. Food Prop.* **2023**, *26(1),* 359-388. DOI: 10.1080/10942912.2022.2157425
5. Mohammed, F.S.; Pehlivan, M.; Sevindik, E.; Akgul, H.; Sevindik, M.; Bozgeyik, I.; Yumrutas, O. Pharmacological properties of edible *Asparagus acutifolius* and *Asparagus officinalis* collected from North Iraq and Turkey (Hatay). *Acta Aliment.* **2021**, *50(1),* 136-143. DOI: 10.1556/066.2020.00204
6. Doğan, M.; Mohammed, F.S.; Uysal, İ.; Mencik, K.; Kına, E.; Pehlivan, M.; Sevindik, M. Total antioxidant status, antimicrobial and antiproliferative potentials of *Viola odorata* (Fragrant Violet). *J. Fac. Pharm. Ankara* **2023**, *47(3),* 784-791. DOI: 10.33483/jfpau.1161440
7. Sevindik, M.; Mohammed, F.S.; Uysal, I. Autism: plants with neuro-psychopharmacotherapeutic potential. *Prospect. Pharm. Sci.* **2023**, *21(3),* 38-48. DOI: 10.56782/pps.143
8. El-Chaghaby, G.A.; Mohammed, F.S.; Rashad, S.; Uysal, I.; Koçer, O.; Lekesiz, Ö.; Sevindik, M. Genus Hypericum: General Properties, Chemical Contents and Biological Activities. *Egypt. J. Botany* **2024**, *64(1),* 1-26. **DOI:** 10.21608/ejbo.2023.217116.2378
9. Colak, S.; Comlekcioglu, N.; Aygan, A.; Kocabas, Y.Z.; Comlekcioglu, U. Phytochemical Properties and Bioactive Potential of Various *Astragalus* spp. from Turkey. *Food Biosci.* **2025**, *64*, Art. No: 105901. DOI: 10.1016/j.fbio.2025.105901
10. Mandour, Y.M.; Refaat, E.; Hassanein, H.D.; Anticancer activity, phytochemical investigation and molecular docking insights of *Citrullus colocynthis* (L.) fruits. *Sci. Rep.* **2023,** 13, Art. No: 20038. DOI: 10.1038/s41598-023-46867-6
11. Cheng, X.; Qin, M.; Chen, R.; Jia, Y.; Zhu, Q.; Chen, G.; Wang, A.; Ling, B.; Rong, W. *Citrullus colocynthis* (L.) Schrad.: A Promising Pharmaceutical Resource for Multiple Diseases. *Molecules* **2023**, *28,* 6221. DOI: 10.3390/molecules28176221
12. Asyaz, S.; Khan, F.U.; Hussain, I.; Khan, M.A.; Khan, I.U. Evaluation of chemical analysis profile of Citrullus colocynthis growing in Southern area of Khyber Pukhtunkhwa, Pakistan. *World Appl. Sci. J.* **2010**, *10*, 402-405.
13. Sevindik, M.; Uysal, İ.; Sevindik, E.; Krupodorova, T.; Koçer, O.; Ünal, O. *Citrullus colocynthis* (Bitter-apple): a comprehensive review on general properties, biological activities, phenolic and chemical contents. *Not. Sci. Biol.* **2024**, *16(4)*, 12202-12202. DOI: 10.55779/nsb16412202

14. Uma, C.; Sekar, K.G. (2014). Phytochemical analysis of a folklore medicinal plant *Citrullus colocynthis* L (bitter apple). *J. Pharmacogn. Phytochem.* **2014**, *2(6),* 195-202.

15. Bauer, A.W.; Kirby, W.M.; Sherris, J.C.; Turck, M. (1966) Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.* **1966**, *45*, 493-496. DOI: 10.1093/ajcp/45.4_ts.493

16. Hindler, J.; Hochstein, L.; Howell, A. Preparation of routine media and reagents used in antimicrobial susceptibility testing. Part 1. McFarland standards, p. 5.19.1-5.19.6. In H. D. Isenberg (ed) *Clinical microbiology procedures handbook* **1992**, vol. 1. American Society for Microbiology, Washington, D.C.

17. Matuschek, E.; Brown, D.F.; Kahlmeter, G. Development of the EUCAST disk diffusion antimicrobial susceptibility testing method and its implementation in routine microbiology laboratories. *Clin. Microbiol. Infect.* **2014**, *20*, 255-266. DOI: 10.1111/1469-0691.12373

18. Baba, H.; Sevindik, M.; Dogan, M.; Akgul, H. Antioxidant, antimicrobial activities and heavy metal contents of some *Myxomycetes*. *Fresenius Environ. Bull.* **2020**, *29(09),* 7840-7846.

19. Erel, O. A novel automated direct measurement method for total antioxidant capacity using a new generation, more stable ABTS radical cation. *Clin. Biochem.* **2004**, *37(4)*, 277-285. DOI: 10.1016/j.clinbiochem.2003.11.015

20. Erel, O. A new automated colorimetric method for measuring total oxidant status. *Clin. Biochem.* **2005**, *38(12),* 1103-1111. DOI: 10.1016/j.clinbiochem.2005.08.008

21. Sevindik, M. The novel biological tests on various extracts of *Cerioporus varius*. *Fresenius Environ. Bull.* **2019**, *28(5*), 3713-3717.

22. Ellman, G.L.; Courtney, K.D.; Andres Jr, V.; Featherstone, R.M. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* **1961**, 7(2), 88-95. DOI: 10.1016/0006-2952(61)90145-9

23. Korkmaz, N.; Koçer, O.; Fathi, S.; Uysal, I.; Sevindik, M. Branched horsetail (*Equisetum ramosissimum*): some biological activities and total phenolic and flavonoid contents. *Prospect. Pharm. Sci.* **2024**, *22(3),* 69-75. DOI: 10.56782/pps.209

24. Sevindik, M.; Bal, C.; Eraslan, E.C.; Uysal, I.; Mohammed, F.S. Medicinal mushrooms: a comprehensive study on their antiviral potential. *Prospect. Pharm. Sci* **2023**, 21(2), 42-56. DOI: 10.56782/pps.141

25. Mohammed, F.S.; Uysal, I.; Sevindik, M. A review on antiviral plants effective against different virus types. *Prospect. Pharm. Sci.* **2023**, *21(2),* 1-21. DOI: 10.56782/pps.128

26. Sevindik, M.; Khassanov, V.T.; Sevindik, E.; Uysal, I.; Mohammed, F.S.. Cornelian Cherry (*Cornus mas* L.): A Comprehensive Review on its Usage Areas, Biological Activities, Mineral, Phenolic and Chemical Contents and Applications. *Appl. Fruit Sci.* **2024**, *66(5),* 2061-2071. DOI: 10.1007/s10341-024-01151-3

27. Marzouk, B.; Marzouk, Z.; Mastouri, M.; Fenina, N.; Aouni, M. Comparative evaluation of the antimicrobial activity of *Citrullus colocynthis* immature fruit and seed organic extracts. *Afr. J. Biotechnol.* **2011**, *10(11),* 2130-2134. DOI: 10.5897/AJB10.1412

28. Gurudeeban, S.; Rajamanickam, E.; Ramanathan, T.; Satyavani, K. Antimicrobial activity of *Citrullus colocynthis* in Gulf of Mannar. *Int. J. Curr. Res.* **2010**, 2(1), 78-81.

29. Tiku, A.R. Antimicrobial compounds and their role in plant defense. In: *Molecular aspects of plant-pathogen interaction,* Eds: Singh, A.; K. Singh, I.K. **2018**, 283-307. DOI: 10.1007/978-981-10-7371-7_13

30. Huang, W.; Wang, Y.; Tian, W.; Cui, X.; Tu, P.; Li, J.; Liu, X. Biosynthesis investigations of terpenoid, alkaloid, and flavonoid antimicrobial agents derived from medicinal plants. *Antibiotics* **2022**, *11(10)*, Art. No: 1380. DOI: 10.3390/antibiotics11101380

31. Gürgen, A.; Sevindik, M. Single and Multi-Objective Optimization of the Red Pine Mushroom *Lactarius deliciosus* (*Agaricomycetes*) Extraction Conditions Using Artificial Intelligence Methods and Biological Activities of Optimized Extracts. *Int. J. Med. Mushrooms* **2025**, 27 (2), 59-73. DOI: 10.1615/IntJMedMushrooms.2024057054

32. Gürgen, A.; Unal, O.; Sevindik, M. Biological activities of the golden chantarelle mushroom *Cantharellus cibarius (Agaricomycetes)* extracts obtained as a result of single and multi-objective optimization studies. *Int. J. Med. Mushrooms* **2024**, *26(12),* 63-74. DOI: 10.1615/IntJMedMushrooms.2024055569

33. Abeyrathne, E.D.N.S.; Nam, K.; Huang, X.; Ahn, D.U. Plant-and animal-based antioxidants' structure, efficacy, mechanisms, and applications: A review. *Antioxidants* **2022**, *11(5),* Art. No: 1025. DOI: 10.3390/antiox11051025

34. Comlekcioglu, N.; Dağlı, F.; Çömlekcioğlu, U.; Aygan, A. *Cornus mas* ve *Rosa canina* Meyvelerinin Antioksidan Kapasitesi ve Bazı Fitokimyasal Özellikleri. *Turkish Journal of Agriculture-Food Science and Technology* **2022**, *10(9),* 1724-1731. DOI: 10.24925/turjaf.v10i9.1724-1731.5434

35. Dumanović, J.; Nepovimova, E.; Natić, M.; Kuča, K.; Jaćević, V. The significance of reactive oxygen species and antioxidant defense system in plants: A concise overview. *Front. Plant Sci.* **2021**, *11*, Art. No: 552969. DOI: 10.3389/fpls.2020.552969

36. Mohammed, F.S.; Sevindik, M.; Uysal, İ.; Sabik, A.E. Quercetin: derivatives, biosynthesis, biological activity, pharmacological and therapeutic effects. *Prospect. Pharm. Sci* **2023**, *21(3),* 49-56. DOI: 10.56782/pps.147

37. Kumar, S.; Kumar, D.; Saroha, K.; Singh, N.; Vashishta, B. Antioxidant and free radical scavenging potential of *Citrullus colocynthis* (L.) Schrad. methanolic fruit extract. *Acta Pharm.* **2008**, *58(2),* 215-220. DOI: 10.2478/v10007-008-0008-1

38. Hussain, A.I.; Rathore, H.A.; Sattar, M.Z.; Chatha,

S.A.; ud din Ahmad, F.; Ahmad, A.; Johns, E.J. Phenolic profile and antioxidant activity of various extracts from *Citrullus colocynthis* (L.) from the Pakistani flora. *Ind. Crops Prod.* **2013**, *45,* 416-422. DOI: 10.1016/j.indcrop.2013.01.002

39. Bourhia, M.; Bouothmany, K.; Bakrim, H.; Hadrach, S.; Salamatullah, A.M.; Alzahrani, A.; Benbacer, L. Chemical profiling, antioxidant, antiproliferative, and antibacterial potentials of chemically characterized extract of *Citrullus colocynthis* L. seeds. Separations **2021**, *8(8),* Art. No: 114. DOI: 10.3390/separations8080114
40. Mohammed, F.S.; Sevindik, M.; Uysal, I.; Sevindik, E.; Akgül, H. A natural material for suppressing the effects of oxidative stress: biological activities of *Alcea kurdica*. *Biol. Bull.* **2022**, *49(Suppl* 2), S59-S66. DOI: 10.1134/S1062359022140102
41. Gürgen, A.; Sevindik, M.; Krupodorova, T.; Uysal, I.; Unal, O. Biological activities of *Hypericum spectabile* extract optimized using artificial neural network combined with genetic algorithm application. *BMC Biotechnol.* **2024**, *24(1),* Art. No: 83. DOI: 10.1186/s12896-024-00914-w
42. Özcandır, A.; Mohammed, F.S.; Sevindik, M.; Aykurt, C.; Selamoglu, Z.; Akgül, H. (2024). Phenolic composition, total antioxidant, antiradical and antimicrobial potential of endemic *Glaucium Alakirensis*. *Sigma J. Eng. Nat. Sci.* **2024**, *42(1),* 42-48. DOI: 10.14744/sigma.2024.00006
43. Seğmenoğlu, M.S.; Sevindik, M. Antioxidant and antimicrobial potentials of functional food *Arum Dioscoridis*. *Sigma J. Eng. Nat. Sci.* **2024**, *42(1),* 116-120. DOI: 10.14744/sigma.2024.00013
44. Sevindik, M.; Gürgen, A.; Krupodorova, T.; Uysal, İ.; Koçer, O. A hybrid artificial neural network and multi-objective genetic algorithm approach to optimize extraction conditions of *Mentha longifolia* and biological activities. *Sci. Rep.* **2024**, *14(1),* Art. No: 31403. DOI: 10.1038/s41598-024-83029-8
45. Sabik, A.E.; Sevindik, M.; Mohammed, F.S.; Dogan, M. A New Natural Source against A549 Lung Cancer Cells *Anthemis cotula* and its Biological Activities and Phenolic Contents. *Pharm. Chem. J.* **2024**, *58(7*), 1134-1140. DOI: 10.1007/s11094-024-03251-6
46. Eraslan, E.C.; Altuntas, D.; Baba, H.; Bal, C.; Akgül, H.; Akata, I.; Sevindik, M. Some biological activities and element contents of ethanol extract of wild edible mushroom *Morchella esculenta*. *Sigma J. Eng. Nat. Sci.* **2021**, *39(1),* 24-28.
47. Mohammed, F.S.; Sevindik, M.; Uysal, İ.; Česko, C.; Koraqi, H. Chemical Composition, Biological Activities, Uses, Nutritional and Mineral Contents of Cumin (*Cuminum cyminum*). *Measurement: Food* **2024**, *14*, Art. No: 100157. DOI: 10.1016/j.meafoo.2024.100157
48. Berwal, M.K.; Ram, C.; Gurjar, P.S.; Gora, J.S.; Kumar, R.; Verma, A.K.; Singh, D.; Basile, B.; Rouphael, Y.; Kumar, P. The bioactive compounds and fatty acid profle of bitter apple seed oil obtained in hot, arid environments. *Horticulturae* **2022**, *8(3),* Art. No: 259. DOI: 10.3390/horticulturae8030259
49. Balasundram, N.; Sundram, K.; Samman, S. Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence, and potential uses. *Food Chem.* **2006**, *99(1*), 191-203. DOI: 10.1016/j.foodchem.2005.07.042
50. Dai, J.; Mumper, R.J. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules* **2010**, *15(10),* 7313-7352. DOI: 10.3390/molecules15107313
51. Chang, C.C.; Yang, M.H.; Wen, H.M.; Chern, J.C. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J. Food Compos. Anal.* **2002**, *10(3),* 178-182. DOI: 10.38212/2224-6614.2748
52. Bejenaru, L.E.; Biță, A.; Mogoșanu, G.D.; Segneanu, A.E.; Radu, A.; Ciocîlteu, M.V.; Bejenaru, C. Polyphenols Investigation and Antioxidant and Anticholinesterase Activities of *Rosmarinus officinalis* L. Species from Southwest Romania Flora. *Molecules* **2024**, *29(18),* Art. No: 4438. DOI: 10.3390/molecules29184438
53. Cartuche, L.; Vallejo, C., Castillo, E.; Cumbicus, N.; Morocho, V. Chemical profiling of *Drimys granadensis* (*Winteraceae*) essential oil, and their antimicrobial, antioxidant, and anticholinesterase properties. *Plants* **2024**, *13(13),* Art. No: 1806. DOI: 10.3390/plants 13131806
54. Shadat, A.A.; Ibrahim, A.Y.; Ezzeldin, E.; Alsaid, M.S. Acetylcholinesterase inhibition and antioxidant activity of some medicinal plants for treating neuro degenarative disease. *Afr. J. Tradit. Complement. Altern. Med.* **2015**, *12(3),* 97-103. DOI: 10.4314/ajtcam.v12i3.12
55. Orhan, I.; Şener, B.; Choudhary, M.I.; Khalid, A. Acetylcholinesterase and butyrylcholinesterase inhibitory activity of some Turkish medicinal plants. *J. Ethnopharmacol.* **2004**, *91(1),* 57-60. DOI: 10.1016/j.jep.2003.11.016
56. Mukherjee, P.K.; Kumar, V.; Mal, M.; Houghton, P.J. Acetylcholinesterase inhibitors from plants. *Phytomedicine* **2007**, *14(4),* 289-300. DOI: 10.1016/j.phymed.2007.02.002